

UDC 593.1
AGRIS L60

<https://doi.org/10.33619/2414-2948/110/05>

**FREE-LIVING PROTOZOA (CILIATES AND TESTATE AMOEBAE)
IN THE FRESHWATERS OF THE NAKHCHIVAN AUTONOMOUS REPUBLIC**

©*Seyidova L.*, Nakhchivan State University,
Nakhchivan, Azerbaijan, leylaseyidova2012@gmail.com

**СВОБОДНОЖИВУЩИЕ ПРОСТЕЙШИЕ (ИНФУЗОРИИ И РАКОВИННЫЕ АМЕБЫ)
В ПРЕСНЫХ ВОДАХ НАХЧЫВАНСКОЙ АВТОНОМНОЙ РЕСПУБЛИКИ**

©*Сеидова Л. М.*, Нахчыванский государственный университет,
г. Нахчыван, Азербайджан, leylaseyidova2012@gmail.com

Abstract. In the period from 2017 to 2022, ciliated protozoa and testate amoebae were studied in freshwater ecosystems of the Nakhchivan Autonomous Republic. 110 species of ciliates and 65 species of amoebae were identified. The species diversity of ciliates in freshwater is almost twice as high as in soil. The differences in species diversity between aquatic and soil amoebae are relatively small. The need for standardization of the methods used by researchers is emphasized. Further studies of these groups of protozoa are necessary to obtain more accurate conclusions about their ecological role, relationships, and their importance in various biological processes.

Аннотация. В период с 2017 по 2022 год в пресноводных экосистемах Нахчыванской Автономной Республики проводилось изучение простейших. Выявлено 110 видов инфузорий и 65 видов амеб. Видовое разнообразие инфузорий в пресной воде почти в два раза выше, чем в почве. Различия в видовом разнообразии между водными и почвенными амебами относительно невелики. Дальнейшие исследования этих групп простейших необходимы для получения более точных выводов об их экологической роли, взаимосвязях и их значении в различных биологических процессах.

Keywords: infusoria, testate amoebae, freshwater, soils, Azerbaijan.

Ключевые слова: инфузории, раковинные амебы, пресные воды, почвы, Азербайджан.

Among other protozoan groups, free-living ciliates and amoebae play a pivotal role in the transformation of organic matter [8, 9].

These unicellular organisms serve as primary consumers and, in turn, as a food source for many small aquatic organisms; they actively contribute to the production and decomposition of organic matter at the primary trophic levels [14, 15].

It has long been established that free-living ciliates exhibit rapid responses to even minor environmental changes, making them valuable tools for bioanalysis at both the cellular and community levels [16].

Despite the recognized importance of studying ciliates and testate amoebae, these two primitive groups—particularly testate amoebae—remain under-researched in Azerbaijan compared to Russia. In contrast, amoebae have been extensively studied by numerous foreign researchers [6, 10, 11, 13, 19].

The earliest information on these protozoan groups in Azerbaijan can be found in the works of Weisig (1940) and Alizade (1939; 1942) [5, 6]. Based on their findings, it was established that prior to the modern era of research, approximately 70 species of ciliates and 28 species of testate amoebae were documented in Azerbaijan's fauna. Currently, the study of ciliates in the region has reached a relatively satisfactory level [3, 4].

It should be mentioned that according to recent estimates, 757 species of ciliates and 265 species of amoebae have been recorded in the waters and soils of Azerbaijan [5].

Building on this data, we conducted a comparative study of the species diversity, distribution, and ecological characteristics of these primitive groups of free-living ciliates and amoebae in the freshwaters of the autonomous republic. The classification system for free-living ciliates followed Lynn (2008), while the taxonomy of testate amoebae was based on the eukaryotic system proposed by an international team of experts [1].

Quantitative analysis of both ciliates and testate amoebae was carried out using the 'non-concentrated' live sample counting method [3].

To determine the species composition of ciliates, the kinetome impregnation techniques using nitrate [7] and silver proteinate [2] were widely employed. Identification of testate amoebae was performed using a scanning electron microscope (JCM 6000, JEOL). The similarity of species diversity between protozoan groups in aquatic biotopes across different regions was evaluated using Bray-Curtis cluster analysis [17].

The species' quantitative ratio was calculated as the ratio of individual species counts to the total number of species (N) in the sample. Based on the results, ciliates and testate amoebae were classified into groups according to Tischler's classification [18]:

1. The group of dominating species — up to n/N 5%.
2. The group of sub-dominating species — up to 2% n/N .
3. The group of presedent species — up to n/N 1% (small).
4. The group of sub-presedent species — less than n/N 1% (random).

All the results were processed through Biodiversity Professional 2 computer.

Results of the study

The studies conducted by us revealed that only 10 species of ciliates and 75 species of testate amoebae were identified in the water bodies of the autonomous republic. The species composition and distribution of these protozoan groups are detailed. The findings indicate that the species composition of amoebae across the three collection sites varies significantly, ranging from a minimum of 31 species to a maximum of 43 species in freshwater habitats.

Additionally, it should be noted that certain groups of free-living ciliates and testate amoebae show a clear preference for either aquatic or terrestrial biotopes. During the study period, the following ciliate species were consistently observed in all freshwater environments: *Anigsteinia salinara*, *Blepharisma tardum*, *B. hyalinum*, *Condylostoma reichii*, *Spirostomum teres*, *Urostyla grandis*, *U. magna*, *Halteria grandinella*, and *Aspidisca steini*.

Among the testate amoebae identified in water bodies, the following species were recorded: *Cyclopyxis eurystoma*, *C. penardi*, *Trigonopyxis arcula*, *Centropyxis aculeata var. oblonga*, *C. minuta*, *C. elongata*, *Diffflugia difficilis*, *D. oblonga*, *D. labiosa*, *Pentagonia azerbaijanica*, *P. compressa*, and others.

Long-term observations of seasonal changes in freshwater ciliate colonies revealed a predominantly dual pattern in their qualitative and quantitative development. In small, temporary reservoirs that often dry up during summer, the maximum abundance of ciliates was typically observed in the spring. However, in autumn, when rainwater refilled these reservoirs and increased their total volume, no significant growth in species diversity or abundance was noted.

The generalized results of assessing the significance of ciliate species in freshwater ecosystems revealed a distinct group of species that are present in water bodies almost year-round, occasionally disappearing from communities only during the winter season. These species include *Oxytricha tenella*, *Euplotes harpa*, *E. balteatus*, *Lacrymaria olor*, *Coleps cucullus*, *Cilodonella spiralis*, *Paramecium caudatum*, *Cyclidium citrullus*, and *Uronema elegans*. Notably, these background species, despite their consistent presence, are typically found in low numbers as single specimens. Their population tends to increase alongside other species during the warmer months.

Regarding the quantitatively dominant species in freshwater communities, it is important to note that the core of dominant and subdominant species often varies significantly between different water bodies. This variation is reflected in the replacement of certain dominant species in one water body by others, which may be subdominant or even rare in neighboring water bodies. Additionally, species dominance is often restricted to specific seasons. For instance, members of the genera *Euplotes* and *Aspidisca*, along with lower-group ciliates such as *Halteria grandinella*, *Heterostrombidium calkinsi*, and *Strombidium conicoides*, exhibit peak quantitative growth in the spring. Conversely, species such as *Oxytricha tenella*, *Lacrymaria kahli*, *Mesodinium acarus*, *Uronema nigricans*, and *Carchesium aselli* become numerically dominant in the autumn.

For freshwater testate amoebae, similar patterns observed in ciliates also apply. Background species in freshwater communities are primarily represented by the genera *Arcella*, *Cyclopyxis*, *Centropyxis*, and *Diffflugia*. Dominant and subdominant groups in these environments include *Arcella hemispherica*, *A. diskoides*, *Cyclopyxis eurystoma*, *Centropyxis aculeata*, and the particularly abundant genus *Diffflugia*.

To summarize the above, it can be noted that the species diversity of free-living ciliates (110 species) and testate amoebae (65 species) in the water bodies of the autonomous republic was found to be notably high.

References:

1. Adl, S. M., Simpson, A. G., Farmer, M. A., Andersen, R. A., Anderson, O. R., Barta, J. R., ... & Taylor, M. F. (2005). The new higher level classification of eukaryotes with emphasis on the taxonomy of protists. *Journal of Eukaryotic Microbiology*, 52(5), 399-451. <https://doi.org/10.1111/j.1550-7408.2005.00053.x>
2. Alekperov, I. Kh. (1992). Novaya modifikatsiya impregnatsii kinetoma infuzorii proteinatom serebra. *Zoologicheskii zhurnal*, 71(2), 130. (In Russian).
3. Alekperov, I. Kh. (2005). Atlas svobodnozhivushchikh infuzorii: (klassy Kinetofragminophora, Colpodea, Oligohymenophora, Polyhymenophora). Baku. (In Russian).
4. Alekperov, I. Kh. (2012). Svobodnozhivushchie infuzorii Azerbaidzhana = Freelifving ciliates of Azerbaijan. Baku. (In Russian).
5. Alekperov, I. Kh., Snegovaya, N. Yu., & Tagirova, E. N. (2017). Kadastr svobodnozhivushchikh infuzorii i rakovinnykh ameb Azerbaidzhana. M.: Tovarishestvo nauchnykh izdaniy KMK, 126 s. (In Russian).
6. Bobrov, A. A. (1999). Ekologo-geograficheskie zakonomernosti rasprostraneniya i struktury soobshchestv rakovinnykh ameb: Protozoa: Testacea: avtoref. dis. ... d-r biol. nauk. Moscow. (In Russian).
7. Chatton, É., & Lwoff, A. (1930). Imprégnation, par diffusion argentine, de l'infaciliature des ciliés marins et d'eau douce, après fixation cytologique et sans dessiccation. *CR Soc. Biol*, 104, 834-836.
8. Foissner, W. (2009). Protist diversity and distribution: some basic considerations. *Protist diversity and geographical distribution*, 1-8. https://doi.org/10.1007/978-90-481-2801-3_1

9. Foissner, W. (2015). Terrestrial and semiterrestrial ciliates (Protozoa, Ciliophora) from Venezuela and Galápagos.
10. Gel'tser, Yu. G., Korganova, G. A., & Alekseev, D. A. (1995). *Opredelitel' pochvoobitayushchikh rakovinykh ameb*. Moscow. (In Russian).
11. Korganova, G. A. (2004). K voprosu o sisteme prosteishikh i taksonomicheskom polozhenii rakovinykh ameb (Rhizopoda, Testacea). *Uspekhi sovremennoi biologii*, 124(5), 443-456. (In Russian).
12. Lynn, D. H. (Ed.). (2008). The ciliated protozoa: characterization, classification, and guide to the literature. https://doi.org/10.1007/978-1-4020-8239-9_17
13. Mazei, Yu. A., & Embulaeva, E. A. (2009). Izmenenie soobshchestv pochvoobitayushchikh rakovinykh ameb vdol' lesostepnogo gradienta v Srednem Povolzh'e. *Aridnye ekosistemy*, 15(37), 13-23. (In Russian).
14. Pavlovskaya, T. V. (1969). Eksperimental'noe issledovanie pitaniya nekotorykh vidov infuzorii Chernogo morya. *Uspekhi protozoologii*, (3), 151-152. (In Russian).
15. Pavlovskaya, T. V. (1973). Vliyanie uslovii pitaniya na skorost' potrebleniya pishchi i vremya generatsii infuzorii. *Zoologicheskii zhurnal*, 52(10), 1451. (In Russian).
16. Railkin, A. I. (2011). Kopirovanie soobshchestv mikroperifitona kak metodicheskii priem pri biotestirovanii. In *Ekologiya svobodnozhivushchikh prosteishikh nazemnykh i vodnykh ekosistem. Tezisy dokladov IV Mezhdunarodnogo simpoziuma. Tol'yatti*, 17-21. (In Russian).
17. Sorensen, T. (1948). A method of establishing groups of equal amplitude in plant sociology based on similarity of species content and its application to analyses of the vegetation on Danish commons. *Biologiske skrifter*, 5, 1-34.
18. Tischler, W. (1955). *Synökologie der Landtiere*.
19. Zharikov, V. V. (1999). *Svobodnozhivushchie infuzorii Volgi: Sostav, dinamika i prostranstvenno-vremennoe raspredelenie v usloviyakh polnogo gidrotekhnicheskogo zaregulirovaniya reki: avtoref. dis. ... d-r biol. nauk*. St. Petersburg. (In Russian).
20. Zharikov, V. V., & Bykova, S. V. (2001). Infuzorii perifitona Saratovskogo vodokhranilishcha. *Fundamental'nye i prikladnye aspekty funktsionirovaniya vodnykh ekosistem*, 263-266. (In Russian).

Список литературы:

1. Adl S. M., Simpson A. G., Farmer M. A., Andersen R. A., Anderson O. R., Barta J. R., Taylor M. F. The new higher level classification of eukaryotes with emphasis on the taxonomy of protists // *Journal of Eukaryotic Microbiology*. 2005. V. 52. №5. P. 399-451. <https://doi.org/10.1111/j.1550-7408.2005.00053.x>
2. Алекперов И. Х. Новая модификация импрегнации кинетома инфузорий протеинатом серебра // *Зоологический журнал*. 1992. Т. 71. №2. С. 130.
3. Алекперов И. Х. Атлас свободноживущих инфузорий: (классы Kinetofragminophora, Colpodea, Oligohymenophora, Polyhymenophora). Баку: Borçali, 2005. 309 с.
4. Алекперов И. Х. Свободноживущие инфузории Азербайджана = Freelifving ciliates of Azerbaijan. Баку: ЭЛМ, 2012. 519 с.
5. Алекперов И. Х., Снеговая Н. Ю., Тагирова Э. Н. Кадастр свободноживущих инфузорий и раковинных амёб Азербайджана. М.: Товарищество научных изданий КМК, 2017. 126 с.
6. Бобров А. А. Эколого-географические закономерности распространения и структуры сообществ раковинных амёб: Protozoa: Testacea: автореф. дис. ... д-р биол. наук. М., 1999. 47 с.

7. Chatton É., Lwoff A. Imprégnation, par diffusion argentine, de l'infraction des ciliés marins et d'eau douce, après fixation cytologique et sans dessiccation // CR Soc. Biol. 1930. V. 104. P. 834-836.
8. Foissner W. Protist diversity and distribution: some basic considerations // Protist diversity and geographical distribution. 2009. P. 1-8. https://doi.org/10.1007/978-90-481-2801-3_1
9. Foissner W. Terrestrial and semiterrestrial ciliates (Protozoa, Ciliophora) from Venezuela and Galápagos. 2015.
10. Гельцер Ю. Г., Корганова Г. А., Алексеев Д. А. Определитель почвообитающих раковинных амёб. М.: Изд-во МГУ, 1995. 83 с.
11. Корганова Г. А. К вопросу о системе простейших и таксономическом положении раковинных амёб (Rhizopoda, Testacea) // Успехи современной биологии. 2004. Т. 124. №5. С. 443-456.
12. Lynn D. H. (ed.). The ciliated protozoa: characterization, classification, and guide to the literature. 2008. https://doi.org/10.1007/978-1-4020-8239-9_17
13. Мазей Ю. А., Ембулаева Е. А. Изменение сообществ почвообитающих раковинных амёб вдоль лесостепного градиента в Среднем Поволжье // Аридные экосистемы. 2009. Т. 15. №37. С. 13-23.
14. Павловская Т. В. Экспериментальное исследование питания некоторых видов инфузорий Черного моря // Успехи протозоологии. 1969. №3. С. 151-152.
15. Павловская Т. В. Влияние условий питания на скорость потребления пищи и время генерации инфузорий // Зоологический журнал. 1973. Т. 52. №10. С. 1451.
16. Раилкин А. И. Копирование сообществ микроперифитона как методический прием при биотестировании // Экология свободноживущих простейших наземных и водных экосистем: Тезисы докладов IV Международного симпозиума. Тольятти. 2011. С. 17-21.
17. Sorensen T. A method of establishing groups of equal amplitude in plant sociology based on similarity of species content and its application to analyses of the vegetation on Danish commons // Biologiske skrifter. 1948. V. 5. P. 1-34.
18. Tischler W. Synökologie der Landtiere. 1955.
19. Жариков В. В. Свободноживущие инфузории Волги: Состав, динамика и пространственно-временное распределение в условиях полного гидротехнического регулирования реки: автореф. дис. ... д-р биол. наук. СПб, 1999. 45 с.
20. Жариков В. В., Быкова С. В. Инфузории перифитона Саратовского водохранилища // Фундаментальные и прикладные аспекты функционирования водных экосистем. 2001. С. 263-266.

Работа поступила
в редакцию 02.12.2024 г.

Принята к публикации
10.12.2024 г.

Ссылка для цитирования:

Seyidova L. Free-Living Protozoa (Ciliates and Testate Amoebae) in the Freshwaters of the Nakhchivan Autonomous Republic // Бюллетень науки и практики. 2025. Т. 11. №1. С. 40-44. <https://doi.org/10.33619/2414-2948/110/05>

Cite as (APA):

Seyidova, L. (2025). Free-Living Protozoa (Ciliates and Testate Amoebae) in the Fresh Waters of the Nakhchivan Autonomous Republic. *Bulletin of Science and Practice*, 11(1), 40-44. <https://doi.org/10.33619/2414-2948/110/05>